arXiv:1303.6760v1 [math.CV] 27 Mar 2013

LANDAU'S THEOREM FOR POLYHARMONIC MAPPINGS

J. CHEN, A. RASILA, AND X. WANG *

Abstract. In this paper, we first investigate coefficient estimates for bounded polyharmonic mappings in the unit disk $\mathbb{D}$. Then, we obtain two versions of Landau's theorem for polyharmonics mapping F, and for the mappings of the type $L(F)$, where L is the differential operator of Abdulhadi, Abu Muhanna and Khuri. Examples and numerical estimates are given.

1. Introduction and preliminaries

A $2p$ ($p \geq 1$) times continuously differentiable complex-valued mapping $F = u+iv$ in a domain $D \subset \mathbb{C}$ is said to be *polyharmonic* (or *p-harmonic*) if F satisfies the polyharmonic equation $\Delta^p F = \Delta(\Delta^{p-1}F) = 0$, where $\Delta := \Delta^1$ is the usual complex Laplacian operator

$$\Delta = 4\frac{\partial^2}{\partial z \partial \overline{z}} := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}.$$

Obviously, for $p = 1$ (resp. $p = 2$), we obtain the usual class of harmonic (resp. biharmonic) mappings. The reader is referred to [8, 9, 22] for discussion on biharmonic and polyharmonic mappings, and [12, 14] for the general theory of harmonic mappings.

The biharmonic equation arises from certain problems in physics, in particular, fluid dynamics and elasticity theory. This equation also has important applications in engineering and biology (cf. [16, 17, 18]). The classes of biharmonic and polyharmonic mappings can be understood as natural generalizations of the class of harmonic mappings. However, the investigation of these mappings in the geometric function theory has been started only recently (cf. [1, 2, 3, 5, 6, 9, 10, 11]).

Let $\mathbb{D}_r = \{z : |z| < r\}$ ($r > 0$), and denote by $\mathbb{D}$ the unit disk $\mathbb{D}_1$. The classical theorem of Landau states that if f is analytic in $\mathbb{D}$, with $f(0) = f'(0) - 1 = 0$ and $|f(z)| < M$ for some $M \geq 1$, then f is univalent in the disk $\mathbb{D}_\rho$ with

$$\rho = \frac{1}{M + \sqrt{M^2 - 1}} > 0.$$

In addition, the range $f(\mathbb{D}_\rho)$ contains a disk of radius $M\rho^2$ (cf. [4]). Recently, there has been a number of articles dealing with Landau's theorem for planar harmonic

---

2000 *Mathematics Subject Classification.* Primary: 30C45; Secondary: 30C20, 30C65.

Key words and phrases. polyharmonic mapping, Landau's theorem.

[*] Corresponding author.

The research was partly supported by NSF of China (No. 11071063) and Hunan Provincial Innovation Foundation for Postgraduate (No. 125000-4242).

mappings, see, for example, [4, 6, 7, 13, 15], and for biharmonic mappings, see [1, 5, 6, 19].

In this paper, we study coefficient estimates for certain classes of polyharmonic mappings. Our goal is to prove Landau type results for these mappings. The main results are Theorems 1, 2 and 3, which are presented in Section 3. Theorem 1 is a generalization of [20, Lemma 2.3], Theorem 2 is a generalization of [8, Theorem 2] and Theorem 3 is a generalization of [5, Theorem 1.1].

2. Preliminaries

If D is a simply connected domain, then it is easy to see that a mapping F is polyharmonic if and only if F has the following representation:

$$F(z)=\sum_{k=1}^{p}|z|^{2(k-1)}G_k(z),$$

where each G_k is a complex-valued harmonic mapping in D for $k\in\{1,\cdots,p\}$ (cf. [8, 22]). It is known (cf. [12, 14]) that the mappings G_k can be expressed as the form

$$G_k=h_k+\overline{g_k}$$

for $k\in\{1,\cdots,p\}$, where all h_k and g_k are analytic in D. In this paper, we consider the polyharmonic mappings in $\mathbb{D}$.

For a polyharmonic mapping F in $\mathbb{D}$, we use the following standard notations:

$$\lambda_F(z)=\min_{0\leq\theta\leq 2\pi}|F_z(z)+e^{-2i\theta}F_{\overline{z}}(z)|=\big||F_z(z)|-|F_{\overline{z}}(z)|\big|,$$

$$\Lambda_F(z)=\max_{0\leq\theta\leq 2\pi}|F_z(z)+e^{-2i\theta}F_{\overline{z}}(z)|=|F_z(z)|+|F_{\overline{z}}(z)|.$$

Thus, for a sense-preserving polyharmonic mapping F, one has $J_F(z)=\Lambda_F(z)\lambda_F(z)$.

We now recall some lemmas, which gives us some coefficient estimates.

Lemma A. [21, Lemma 2.1] *Suppose that $f=h+\overline{g}$ is a harmonic mapping of $\mathbb{D}$ with $h(z)=\sum_{n=1}^{\infty}a_nz^n$ and $g(z)=\sum_{n=1}^{\infty}b_nz^n$ for $z\in\mathbb{D}$. If $J_F(0)=1$ and $|f(z)|<M$, then*

$$|a_n|,\ |b_n|\leq\sqrt{M^2-1},\quad n=2,3,\cdots,$$
$$|a_n|+|b_n|\leq\sqrt{2M^2-2},\quad n=2,3,\cdots,$$

and

$$\lambda_F(0)\geq\lambda(M)=\begin{cases}\dfrac{\sqrt{2}}{\sqrt{M^2-1}+\sqrt{M^2+1}} & \text{if } 1\leq M\leq\frac{\pi}{2\sqrt[4]{2\pi^2-16}},\\[2ex] \dfrac{\pi}{4M} & \text{if } M>\frac{\pi}{2\sqrt[4]{2\pi^2-16}}.\end{cases}$$

Lemma B. [8, Lemma 1] *Let $f=h+\overline{g}$ be a harmonic mapping of $\mathbb{D}$ such that $|f(z)|<M$ with $h(z)=\sum_{n=0}^{\infty}a_nz^n$ and $g(z)=\sum_{n=1}^{\infty}b_nz^n$. Then $|a_0|\leq M$ and for any $n\geq1$,*

$$|a_n|+|b_n|\leq\frac{4M}{\pi}. \tag{1}$$

The estimate (1) *is sharp. The extremal functions are* $f(z) \equiv M$ *or*

$$f_n(z) = \frac{2M\alpha}{\pi} \arg\left(\frac{1+\beta z^n}{1-\beta z^n}\right),$$

where $|\alpha| = |\beta| = 1$.

Lemma C. [20, Lemma 2.3] *Suppose that* $f = h + \overline{g}$ *is a harmonic mapping of the unit disk* $\mathbb{D}$ *with* $h(z) = \sum_{n=0}^{\infty} a_n z^n$ *and* $g(z) = \sum_{n=1}^{\infty} b_n z^n$.

(*I*) *If* f *satisfies* $|f(z)| \leq M$ *for all* $z \in \mathbb{D}$, *then*

$$\sum_{n=2}^{\infty} (|a_n|^2 + |b_n|^2) \leq M^2 - (|a_1|^2 + |b_1|^2).$$

(*II*) *If* f *satisfies* $|f(z)| \leq M$ *for all* $z \in \mathbb{D}$ *and* $|J_F(0)| = 1$, *then*

$$\left(\sum_{n=2}^{\infty} (|a_n|^2 + |b_n|^2)\right)^{\frac{1}{2}} \leq \sqrt{M^4 - 1} \cdot \lambda_F(0).$$

(*III*) *If* f *satisfies* $|f(z)| \leq M$ *for all* $z \in \mathbb{D}$ *and* $\lambda_F(0) = 1$, *then*

$$\left(\sum_{n=2}^{\infty} (|a_n|^2 + |b_n|^2)\right)^{\frac{1}{2}} \leq \sqrt{2(M^2 - 1)}.$$

In [3], the authors considered the following differential operator L defined on the class of complex-valued C^1 functions:

$$L = z\frac{\partial}{\partial z} - \overline{z}\frac{\partial}{\partial \overline{z}}.$$

Clearly, L is a complex linear operator, and L satisfies the usual product rule:

$$L(af + bg) = aL(f) + bL(g) \text{ and } L(fg) = fL(g) + gL(f),$$

where a, b are complex constants, and f, g are C^1 functions. In addition, the operator L has a number of interesting properties. For instance, it is easy to see that the operator L preserves both harmonicity and biharmonicity. Other basic properties of this operator are presented in [3].

Theorem D. [3, Corollary 1(3)] *Let* F *be a univalent biharmonic function in* $\mathbb{D}$. *If* F *is convex and* $L(F)$ *is univalent, then* $L(F)$ *is starlike.*

It is also important to recall that (see [3, Corollary 1(3)]) the operator $L(F)$ for polyharmonic mappings behaves much like $zf'(z)$ for analytic functions. From Theorem D, we see that it is meaningful to consider Landau's theorem for polyharmonic mappings of the form $L(F)$, where F belongs to the class of polyharmonic mappings.

3. Main results

We begin this section with some coefficient estimates for polyharmonic mappings.

Theorem 1. *Suppose that F is a polyharmonic mapping of the form*

$$F(z)=a_0+\sum_{k=1}^{p}|z|^{2(k-1)}(h_k(z)+\overline{g_k(z)})=a_0+\sum_{k=1}^{p}|z|^{2(k-1)}\sum_{n=1}^{\infty}(a_{n,k}z^n+\overline{b_{n,k}}\overline{z^n}), \tag{2}$$

and all its non-zero coefficients a_{n,k_1}, a_{n,k_2} and b_{n,k_3}, b_{n,k_4} satisfy the condition:

$$\left|\arg\frac{a_{n,k_1}}{a_{n,k_2}}\right|\leq\frac{\pi}{2}\ \text{and}\ \left|\arg\frac{b_{n,k_3}}{b_{n,k_4}}\right|\leq\frac{\pi}{2}. \tag{3}$$

(I) If F satisfies $|F(z)|\leq M$ for all $z\in\mathbb{D}$, then

$$|a_0|^2+\sum_{k=1}^{p}\sum_{n=1}^{\infty}(|a_{n,k}|^2+|b_{n.k}|^2)\leq M^2, \tag{4}$$

and $|a_{n,k}|+|b_{n,k}|\leq\sqrt{2}M$ for all coefficients.

(II) If F satisfies $|F(z)|\leq M$ for all $z\in\mathbb{D}$ and $F(0)=|J_F(0)|-1=0$, then

$$\sqrt{\sum_{k=1}^{p}\sum_{n=1}^{\infty}(|a_{n,k}|+|b_{n.k}|)^2-(|a_{1,1}|+|b_{1,1}|)^2}\leq\sqrt{M^4-1}\cdot\lambda_F(0), \tag{5}$$

and $|a_{n,k}|+|b_{n,k}|\leq T_1(M):=\min\left\{\sqrt{2M^2-2},\ \sqrt{M^4-1}\cdot\lambda_F(0)\right\}$ for all $(n,k)\neq(1,1)$, where

$$\lambda_F(0)\geq\lambda_0(M):=\frac{\sqrt{2}}{\sqrt{M^2-1}+\sqrt{M^2+1}}. \tag{6}$$

(III) If F satisfies $|F(z)|\leq M$ for all $z\in\mathbb{D}$ and $F(0)=\lambda_F(0)-1=0$, then

$$\sqrt{\sum_{k=1}^{p}\sum_{n=1}^{\infty}(|a_{n,k}|+|b_{n.k}|)^2-(|a_{1,1}|+|b_{1,1}|)^2}\leq\sqrt{2M^2-2}, \tag{7}$$

and $|a_{n,k}|+|b_{n,k}|\leq\sqrt{2M^2-2}$ for all $(n,k)\neq(1,1)$.

Proof. We first prove the inequality (4). A standard argument using Parseval's identity and the hypothesis that $|F(re^{i\theta})|\leq M$, gives

$$\begin{aligned}&\frac{1}{2\pi}\int_0^{2\pi}|F(re^{i\theta})|^2d\theta\\ =\ &|a_0|^2+\sum_{k=1}^{p}r^{4(k-1)}\sum_{n=1}^{\infty}(|a_{n,k}|^2+|b_{n,k}|^2)r^{2n}\\ &+2\mathrm{Re}\sum_{1\leq k_1<k_2\leq p}r^{2(k_1+k_2-2)}\sum_{n=1}^{\infty}(a_{n,k_1}\overline{a_{n,k_2}}+b_{n,k_1}\overline{b_{n,k_2}})r^{2n}\\ \leq\ &M^2.\end{aligned}$$

Since F satisfies the condition (3), it follows that $\mathrm{Re}(a_{n,k_1}\overline{a_{n,k_2}} + b_{n,k_1}\overline{b_{n,k_2}}) \geq 0$ for any k_1, $k_2 \in \{1, \cdots, p\}$. Let $r \longrightarrow 1^-$. We obtain that

$$|a_0|^2 + \sum_{k=1}^{p}\sum_{n=1}^{\infty}(|a_{n,k}|^2 + |b_{n.k}|^2) \leq M^2,$$

which proves that the inequality (4) holds. By Cauchy's inequality, for any coefficient $a_{n,k}$, $b_{n,k}$, we get

$$(|a_{n,k}| + |b_{n,k}|)^2 \leq 2(|a_{n,k}|^2 + |b_{n,k}|^2) \leq 2M^2.$$

That is $|a_{n,k}| + |b_{n,k}| \leq \sqrt{2}M$.

Then, we prove the inequality (5). We have from (4) that

$$\begin{aligned}
&\left(\sum_{k=1}^{p}\sum_{n=2}^{\infty}(|a_{n,k}| + |b_{n.k}|)^2 + \sum_{k=2}^{p}(|a_{1,k}| + |b_{1.k}|)^2\right)^{\frac{1}{2}} \\
\leq &\left(\sum_{k=1}^{p}\sum_{n=2}^{\infty}2(|a_{n,k}|^2 + |b_{n.k}|^2) + \sum_{k=2}^{p}2(|a_{1,k}|^2 + |b_{1.k}|^2)\right)^{\frac{1}{2}} \\
\leq &\big(2M^2 - 2(|a_{1,1}|^2 + |b_{1,1}|^2)\big)^{\frac{1}{2}}.
\end{aligned} \tag{8}$$

It follows from (8) that $|a_{n,k}| + |b_{n.k}| \leq \sqrt{2M^2 - 2}$ for $(n, k) \neq (1, 1)$. Then if $J_F(0) = 1$, we get $|a_{1,1}| = \sqrt{1 + |b_{1,1}|^2} \geq 1$. Thus we deduce from (4) that $|b_{1,1}| \leq \sqrt{\frac{M^2-1}{2}}$, $M \geq 1$. Hence

$$\sqrt{2M^2 - 2(|a_{1,1}|^2 + |b_{1,1}|^2)} = \sqrt{2M^2 - 2(1 + 2|b_{1,1}|^2)}\Big(|b_{1,1}| + \sqrt{1 + |b_{1,1}|^2}\Big) \cdot \lambda_F(0).$$

As in [20], by a straightforward calculation, we get

$$\sqrt{2M^2 - 2(1 + 2|b_{1,1}|^2)}\Big(|b_{1,1}| + \sqrt{1 + |b_{1,1}|^2}\Big) \leq \sqrt{M^4 - 1},$$

which proves that the inequality (5) holds, $|a_{n,k}| + |b_{n.k}| \leq \sqrt{M^4 - 1} \cdot \lambda_F(0)$ for $(n, k) \neq (1, 1)$ and

$$\lambda_F(0) = \big||a_{1,1}| - |b_{1,1}|\big| = \frac{1}{\sqrt{1 + |b_{1,1}|^2} + |b_{1,1}|} \geq \frac{\sqrt{2}}{\sqrt{M^2 - 1} + \sqrt{M^2 + 1}}.$$

If $J_F(0) = -1$, we get $|b_{1,1}| = \sqrt{1 + |a_{1,1}|^2} \geq 1$. Thus we deduce from (4) that $|a_{1,1}| \leq \sqrt{\frac{M^2-1}{2}}$, $M \geq 1$. Hence

$$\sqrt{2M^2 - 2(|a_{1,1}|^2 + |b_{1,1}|^2)} = \sqrt{2M^2 - 2(1 + 2|a_{1,1}|^2)}\Big(|a_{1,1}| + \sqrt{1 + |a_{1,1}|^2}\Big) \cdot \lambda_F(0).$$

As in the previous case, we get that

$$\sqrt{2M^2 - 2(1 + 2|a_{1,1}|^2)}\Big(|a_{1,1}| + \sqrt{1 + |a_{1,1}|^2}\Big) \leq \sqrt{M^4 - 1},$$

which proves that the inequality (5) holds, $|a_{n,k}| + |b_{n.k}| \leq \sqrt{M^4-1} \cdot \lambda_F(0)$ for $(n,k) \neq (1,1)$ and

$$\lambda_F(0) = \big||a_{1,1}| - |b_{1,1}|\big| = \frac{1}{\sqrt{1+|a_{1,1}|^2}+|a_{1,1}|} \geq \frac{\sqrt{2}}{\sqrt{M^2-1}+\sqrt{M^2+1}}.$$

Next, we prove the inequality (7). We have from $\lambda_F(0) = \big||a_{1,1}| - |b_{1,1}|\big| = 1$ that $|a_{1,1}|^2 + |b_{1,1}|^2 \geq 1$. Hence the inequality (7) follows from (8). □

From (8), we obtain the following two corollaries.

Corollary 1. *Suppose that F is a polyharmonic mapping of the form* (2) *and all its non-zero coefficients a_{n,k_1}, a_{n,k_2} and b_{n,k_3}, b_{n,k_4} satisfy* (3).

If F satisfies $|F(z)| \leq M$ for all $z \in \mathbb{D}$, then for all $n \in N^*$,

$$\sum_{k=1}^{p} |a_{n,k}|,\ \sum_{k=1}^{p} |b_{n,k}| \leq \sqrt{p}M \text{ and } \sum_{k=1}^{p}(|a_{n,k}| + |b_{n,k}|) \leq \sqrt{2p}M.$$

In addition, if $F(0)=0$, $\lambda_F(0)=1$ or $|J_F(0)|=1$, then $|a_{n,k}|$, $|b_{n,k}| \leq \sqrt{M^2-1}$ and for $n \in N^* \setminus \{1\}$

$$\sum_{k=1}^{p} |a_{n,k}|,\ \sum_{k=1}^{p} |b_{n,k}| \leq \sqrt{p(M^2-1)} \text{ and } \sum_{k=1}^{p}(|a_{n,k}| + |b_{n,k}|) \leq \sqrt{2p(M^2-1)}.$$

Corollary 2. *Suppose that F is a polyharmonic mapping of the form* (2)*, all its non-zero coefficients a_{n,k_1}, a_{n,k_2} and b_{n,k_3}, b_{n,k_4} satisfy* (3)*, $F(0)=0$, $|F(z)| \leq 1$ on $\mathbb{D}$.*

(I) If $J_F(0) = 1$, then $F(z) = \alpha z$, where $|\alpha| = 1$. And if $J_F(0) = -1$, then $F(z) = \beta\overline{z}$, where $|\beta| = 1$.

(II) If $\lambda_F(0) = 1$, then $F(z) = \gamma z$ or $F(z) = \gamma\overline{z}$, where $|\gamma| = 1$.

Example 1. *Let $\beta = \sqrt{\alpha} = e^{i\pi/n}$ and*

$$\begin{aligned} f_n(z) &= \frac{1}{2\pi i}\sum_{k=0}^{n-1} \alpha^k \left\{\log\frac{z-\beta^{2k+1}}{z-\beta^{2k-1}} - \overline{\log\frac{z-\beta^{2k+1}}{z-\beta^{2k-1}}}\right\} \\ &= \sum_{m=1}^{\infty}\left(a_m z^m + \overline{b_m}\overline{z^m}\right), \end{aligned}$$

where

$$a_m = \begin{cases} \dfrac{n}{\pi m}\sin\dfrac{\pi m}{n}, & m = 1,\ n+1,\ 2n+1, \cdots, \\ 0, & \text{otherwise}, \end{cases}$$

$$b_m = \begin{cases} \dfrac{n}{\pi m}\sin\dfrac{\pi m}{n}, & m = n-1,\ 2n-1, \cdots, \\ 0, & \text{otherwise}, \end{cases}$$

for $n = 3, 4, \cdots$. Then f_n is univalent in $\mathbb{D}$, and maps $\mathbb{D}$ onto the region inside the regular n-gon with vertices 1, $\alpha, \cdots, \alpha^{n-1}$ (cf. [14, p. 59]*). Let $F_0(z) = f_3(z) +$*



FIGURE 1. The images of $\mathbb{D}$ under f_3 (left) and F_0 (right). See Example 1.

$17i|z|^2f_3(z)$. *Then* $|F_0| < 18$ *in* $\mathbb{D}$, *and* F_0 *satisfies the condition* (3). *Therefore, by* (4) *we obtain*

$$\sum_{k=1}^{p}\sum_{n=1}^{\infty}(|a_{n,k}|^2+|b_{n.k}|^2)\leq 324,$$

and $|a_{n,k}|+|b_{n,k}|\leq 18\sqrt{2}$ *for all coefficients. See Figure 1 for the images of* $\mathbb{D}$ *under* f_3 *and* F_0.

Now, we establish a version of Landau's theorem for polyharmonic mappings.

Theorem 2. *Suppose that* F *is a polyharmonic mapping of the form* (2) *and all its non-zero coefficients* a_{n,k_1}, a_{n,k_2} *and* b_{n,k_3}, b_{n,k_4} *satisfy* (3).

If $|F(z)|\leq M$ *in* $\mathbb{D}$ *for some* $M>1$ *and* $F(0)=|J_F(0)|-1=0$, *then* F *is univalent in the disk* $\mathbb{D}_{r_1}$ *and* $F(\mathbb{D}_{r_1})$ *contains a univalent disk* $\mathbb{D}_{\rho_1}$, *where* $r_1(M,p)$ *is the least positive root of the following equation:*

$$1-\sqrt{M^4-1}\left(\frac{2r-r^2}{(1-r)^2}+\sum_{k=1}^{p-1}\frac{r^{2k}}{(1-r)^2}+2\sum_{k=1}^{p-1}\frac{kr^{2k}}{1-r}\right)=0,$$

and

$$\rho_1=\lambda_0(M)r_1\left(1-\sqrt{M^4-1}\frac{r_1}{1-r_1}-\sqrt{M^4-1}\sum_{k=1}^{p-1}\frac{2r_1^{2k}}{1-r_1}\right).$$

In particular, if $|F(z)|\leq 1$ *for all* $z\in\mathbb{D}$ *and* $F(0)=|J_F(0)|-1=0$, *then* F *is univalent and maps* $\mathbb{D}$ *onto* $\mathbb{D}$.

Proof. Since $F_z(z)$ and $F_{\overline{z}}(z)$ can be written as

$$F_z(z)=\sum_{k=1}^{p}|z|^{2(k-1)}h_k'(z)+\sum_{k=2}^{p}(k-1)\overline{z}|z|^{2(k-2)}\big(h_k(z)+\overline{g_k(z)}\big),$$

$$F_{\overline{z}}(z)=\sum_{k=1}^{p}|z|^{2(k-1)}\overline{g_k'(z)}+\sum_{k=2}^{p}(k-1)z|z|^{2(k-2)}\big(h_k(z)+\overline{g_k(z)}\big),$$

then for any $z_1\neq z_2$, where z_1, $z_2\in\mathbb{D}_r$ and $r\in(0,1)$ is a constant, we have

$$\begin{aligned}\big|F(z_1)-F(z_2)\big| &= \left|\int_{[z_1,z_2]}F_z(z)dz+F_{\overline{z}}(z)d\overline{z}\right|\\ &\geq \left|\int_{[z_1,z_2]}F_z(0)dz+F_{\overline{z}}(0)d\overline{z}\right|\\ &\quad -\left|\int_{[z_1,z_2]}\big(F_z(z)-F_z(0)\big)dz+\big(F_{\overline{z}}(z)-F_{\overline{z}}(0)\big)d\overline{z}\right|\\ &\geq J_1-J_2-J_3-J_4,\end{aligned}$$

where

$$\begin{aligned}J_1 &:= \left|\int_{[z_1,z_2]}h_1'(0)\,dz+\overline{g_1'(0)}d\overline{z}\right|,\\ J_2 &:= \left|\int_{[z_1,z_2]}\big(h_1'(z)-h_1'(0)\big)dz+\big(\overline{g_1'(z)}-\overline{g_1'(0)}\big)d\overline{z}\right|,\\ J_3 &:= \left|\int_{[z_1,z_2]}\sum_{k=1}^{p-1}|z|^{2k}h_{k+1}'(z)dz+\sum_{k=1}^{p-1}|z|^{2k}\overline{g_{k+1}'(z)}d\overline{z}\right|,\\ J_4 &:= \left|\int_{[z_1,z_2]}\sum_{k=1}^{p-1}k|z|^{2(k-1)}\big(h_{k+1}(z)+\overline{g_{k+1}(z)}\big)(\overline{z}dz+zd\overline{z})\right|.\end{aligned}$$

First, we have the estimate for J_1,

$$J_1\geq\int_{[z_1,z_2]}\lambda_F(0)|dz|=\lambda_F(0)|z_1-z_2|.$$

Next, by Theorem 1 (II), we obtain $|a_{n,k}|+|b_{n,k}|\leq\sqrt{M^4-1}\cdot\lambda_F(0)$ for $(n,k)\neq(1,1)$. It follows that

$$\begin{aligned}J_2 &\leq \int_{[z_1,z_2]}\big|h_1'(z)-h_1'(0)\big||dz|+\big|\overline{g_1'(z)}-\overline{g_1'(0)}\big||d\overline{z}|\\ &\leq |z_1-z_2|\sum_{n=2}^{\infty}n(|a_{n,1}|+|b_{n,1}|)r^{n-1}\\ &\leq |z_1-z_2|\sqrt{M^4-1}\cdot\lambda_F(0)\frac{2r-r^2}{(1-r)^2},\end{aligned}$$

and

$$\begin{aligned}
J_3 &\leq \int_{[z_1,z_2]} \sum_{k=1}^{p-1} |z|^{2k} \big|h'_{k+1}(z)\big||dz| + \sum_{k=1}^{p-1} |z|^{2k} \big|\overline{g'_{k+1}(z)}\big||d\overline{z}| \\
&\leq |z_1 - z_2| \sum_{k=1}^{p-1} r^{2k} \sum_{n=1}^{\infty} n(|a_{n,k+1}| + |b_{n,k+1}|) r^{n-1} \\
&\leq |z_1 - z_2| \sqrt{M^4 - 1} \cdot \lambda_F(0) \sum_{k=1}^{p-1} \frac{r^{2k}}{(1-r)^2}.
\end{aligned}$$

Finally,

$$\begin{aligned}
J_4 &= \left| \int_{[z_1,z_2]} \sum_{k=1}^{p-1} k|z|^{2(k-1)} \big(h_{k+1}(z) + \overline{g_{k+1}(z)}\big)(\overline{z}dz + zd\overline{z}) \right| \\
&\leq \left| \int_{[z_1,z_2]} 2 \sum_{k=1}^{p-1} kr^{2(k-1)} \sum_{n=1}^{\infty} (|a_{n,k+1}| + |b_{n,k+1}|)\, r^{n+1} |dz| \right| \\
&\leq 2|z_1 - z_2| \sum_{k=1}^{p-1} kr^{2(k-1)} \sum_{n=1}^{\infty} (|a_{n,k+1}| + |b_{n,k+1}|)\, r^{n+1} \\
&\leq 2|z_1 - z_2| \sqrt{M^4 - 1} \cdot \lambda_F(0) \sum_{k=1}^{p-1} \frac{kr^{2k}}{1-r}.
\end{aligned}$$

Using these estimates, we obtain

$$\big|F(z_1) - F(z_2)\big| \geq J_1 - J_2 - J_3 - J_4 \geq \lambda_F(0)|z_1 - z_2|\psi(r),$$

where

$$\psi(r) = 1 - \sqrt{M^4 - 1} \left(\frac{2r - r^2}{(1-r)^2} + \sum_{k=1}^{p-1} \frac{r^{2k}}{(1-r)^2} + 2\sum_{k=1}^{p-1} \frac{kr^{2k}}{1-r} \right).$$

It is easy to see that the function $\psi(r)$ is strictly decreasing for $r \in (0,1)$ and $M > 1$,

$$\lim_{r\to 0+} \psi(r) = 1 \text{ and } \lim_{r\to 1^-} \psi(r) = -\infty.$$

Hence there exists a unique $r_1 \in (0,1)$ satisfying $\psi(r_1) = 0$. This implies that F is univalent in $\mathbb{D}_{r_1}$.

By Theorem 1 (II), we see $|a_{n,k}|+|b_{n,k}|\leq\sqrt{M^4-1}\cdot\lambda_F(0)$ for all $(n,k)\neq(1,1)$. Therefore, by (6), for any w in $\{w:\ |w|=r_1\}$, we obtain

$$\begin{aligned}
\big|F(w)-F(0)\big| &= \left|\int_{[0,w]}F_z(z)dz+F_{\overline{z}}(z)d\overline{z}\right|\\
&\geq \lambda_F(0)r_1-\sum_{n=2}^{\infty}(|a_{n,1}|+|b_{n,1}|)r_1^n-2\sum_{k=1}^{p-1}r_1^{2k}\sum_{n=1}^{\infty}(|a_{n,k+1}|+|b_{n,k+1}|)r_1^n\\
&\geq \lambda_F(0)r_1\left(1-\sqrt{M^4-1}\frac{r_1}{1-r_1}-\sqrt{M^4-1}\sum_{k=1}^{p-1}\frac{2r_1^{2k}}{1-r_1}\right)\\
&\geq \lambda_0(M)r_1\left(1-\sqrt{M^4-1}\frac{r_1}{1-r_1}-\sqrt{M^4-1}\sum_{k=1}^{p-1}\frac{2r_1^{2k}}{1-r_1}\right):=\rho_1.
\end{aligned}$$

We also have

$$\rho_1>\lambda_0(M)r_1\left[1-\sqrt{M^4-1}\left(\frac{2r_1-r_1^2}{(1-r_1)^2}+\sum_{k=1}^{p-1}\frac{r_1^{2k}}{(1-r_1)^2}+2\sum_{k=1}^{p-1}\frac{kr_1^{2k}}{1-r_1}\right)\right]=0.$$

If $|F(z)|\leq 1$ for all $z\in\mathbb{D}$ and $F(0)=|J_F(0)|-1=0$, then by Corollary 2 (I), we have F is univalent and maps $\mathbb{D}$ onto $\mathbb{D}$. The proof of the theorem is complete. $\square$

Corollary 3. *Suppose that F is a polyharmonic mapping of the form* (2) *and all its non-zero coefficients a_{n,k_1}, a_{n,k_2} and b_{n,k_3}, b_{n,k_4} satisfy* (3).

If $|F(z)|\leq M$ in $\mathbb{D}$ for some $M>1$ and $F(0)=\lambda_F(0)-1=0$, then F is univalent in the disk $\mathbb{D}_{r_2}$ and $F(\mathbb{D}_{r_2})$ contains a univalent disk $\mathbb{D}_{\rho_2}$, where $r_2(M,p)$ is the least positive root of the following equation:

$$1-\sqrt{2M^2-2}\left(\frac{2r-r^2}{(1-r)^2}+\sum_{k=1}^{p-1}\frac{r^{2k}}{(1-r)^2}+2\sum_{k=1}^{p-1}\frac{kr^{2k}}{1-r}\right)=0,$$

and

$$\rho_2=r_2\left(1-\sqrt{2M^2-2}\frac{r_2}{1-r_2}-\sqrt{2M^2-2}\sum_{k=1}^{p-1}\frac{2r_2^{2k}}{1-r_2}\right).$$

In particular, if $|F(z)|\leq 1$ for all $z\in\mathbb{D}$ and $F(0)=\lambda_F(0)-1=0$, then F is univalent and maps $\mathbb{D}$ onto $\mathbb{D}$.

Proof. The proof of this result is similar to Theorem 1, where $|a_{n,k}|+|b_{n,k}|\leq\sqrt{2M^2-2}$ and $\lambda_F(0)=1$ is used instead of $|a_{n,k}|+|b_{n,k}|\leq\sqrt{M^4-1}\cdot\lambda_F(0)$ for all $(n,k)\neq(1,1)$, and we omit it. $\square$

Example 2. *Let $F_1(z)=K(z)+|z^2|G(z)=\frac{2\pi}{3\sqrt{3}}f_3(z)+\frac{34\pi i}{3\sqrt{3}}|z|^2f_3(z)$ for $z\in\mathbb{D}$. Then $F_1(0)=J_{F_1}(0)-1=\lambda_{F_1}(0)-1=0$, $|F_1(z)|<M_1:=4\sqrt{3}\pi$ and both $|K|$ and $|G|$ are bounded by $M_2:=\frac{34\pi}{3\sqrt{3}}$. Then, by Corollary 3, F_1 is univalent in the*

disk $\mathbb{D}_{r_3}$ *and* $F(\mathbb{D}_{r_3})$ *contains a univalent disk* $\mathbb{D}_{\rho_3}$*, where* $r_3 \approx 0.01552$ *is the least positive root of the following equation:*

$$1-\sqrt{2M_1^2-2}\left(\frac{2r}{(1-r)^2}+2\frac{r^2}{1-r}\right)=0,$$

and

$$\rho_3=r_3\left(1-\sqrt{2M_1^2-2}\cdot\frac{r_3+2r_3^2}{1-r_3}\right)\approx 0.00776,$$

while by [8, Theorem 2]*, we see that* F_1 *is univalent in the disk* $\mathbb{D}_{r_4}$ *and* $F_1(\mathbb{D}_{r_4})$ *contains a univalent disk* $\mathbb{D}_{\rho_4}$*, where* $r_4 \approx 0.00041$ *is the least positive root of the following equation:*

$$\frac{\pi}{4M_2}-\frac{4M_2r(2-r)}{\pi(1-r)^2}-\frac{4M_2r^2}{\pi(1-r)^2}-2M_2r=0,$$

and

$$\rho_4=r_4\left(\frac{\pi}{4M_2}-\frac{4M_2r_4}{\pi(1-r_4)}-\frac{4M_2r_4^2}{\pi(1-r_4)}\right)\approx 1.12385\times 10^{-5}.$$

Corollary 4. *Suppose that* F *is a polyharmonic mapping of the form*

$$F(z)=\sum_{k=1}^{p}\lambda_k|z|^{2(k-1)}G(z)=\sum_{k=1}^{p}|z|^{2(k-1)}\sum_{n=1}^{\infty}(a_nz^n+\overline{b_n}\overline{z^n}),$$

where $G(z)$ *is nonconstant,* $\sum_{k=1}^{p}\lambda_k=1$ *and* $\lambda_k\in[0,1]$ *for all* $k\in\{1,\cdots,p\}$*. If* $|F(z)|\le M$ *in* $\mathbb{D}$ *for some* $M>1$ *and* $\lambda_F(0)=1$*, then* F *is univalent in the disk* $\mathbb{D}_{r_5}$*, and* $F(\mathbb{D}_{r_5})$ *contains a univalent disk* $\mathbb{D}_{\rho_5}$*, where* $r_5(M,p)$ *is the least positive root of the following equation:*

$$1-T(M)\left(\frac{2r-r^2}{(1-r)^2}+\sum_{k=1}^{p-1}\frac{r^{2k}}{(1-r)^2}+2\sum_{k=1}^{p-1}\frac{kr^{2k}}{1-r}\right)=0,$$

and

$$\rho_5=r_5\left(1-T(M)\frac{r_5}{1-r_5}-T(M)\sum_{k=1}^{p-1}\frac{2r_5^{2k}}{1-r_5}\right),$$

where $T(M):=\min\left\{\sqrt{2M^2-2},\ \frac{4M}{\pi}\right\}$.

Proof. Suppose that $|F(z)|\le M$ for all $z\in\mathbb{D}$ and $\sum_{k=1}^{p}\lambda_k=1$. Then

$$\lim_{r\to 1^-}|F(re^{i\theta})|=\lim_{r\to 1^-}|G(re^{i\theta})|\le M.$$

Since $G(z)$ is harmonic and nonconstant, by the maximum principle, we have $|G(z)|<M$ in $\mathbb{D}$. Lemma B and Theorem 1 (III) implies that

$$|a_n|+|b_n|\le T(M)=\min\left\{\sqrt{2M^2-2},\frac{4}{\pi}M\right\}$$

for all $n\ge 1$. By Corollary 3, we easily obtain the results. ☐

Now, we consider Landau's theorem for the mapping $L(F)$, where F is polyharmonic.

Theorem 3. *Suppose that F is a polyharmonic mapping of the form (2) and all its non-zero coefficients a_{n,k_1}, a_{n,k_2} and b_{n,k_3}, b_{n,k_4} satisfy (3).*

If $|F(z)| \leq M$ in $\mathbb{D}$ for some $M > 1$ and $F(0) = |J_F(0)| - 1 = 0$, then $L(F)$ is univalent in the disk $\mathbb{D}_{r_6}$ and $F(\mathbb{D}_{r_6})$ contains a univalent disk $\mathbb{D}_{\rho_6}$, where $r_6(M,p)$ is the least positive root of the following equation:

$$1 - \sqrt{M^4-1}\left(\frac{2r-r^2}{(1-r)^2} + \sum_{k=1}^{p}\frac{2r^{2k-1}}{(1-r)^3} + \sum_{k=2}^{p}\frac{(2k-1)r^{2(k-1)}}{(1-r)^2}\right) = 0,$$

and

$$\rho_6 = \lambda_0(M) r_6 \left(1 - \sqrt{M^4-1}\frac{2r_6 - r_6^2}{(1-r_6)^2} - \sqrt{M^4-1}\sum_{k=2}^{p}\frac{r_6^{2(k-1)}}{(1-r_6)^2}\right).$$

In particular, if $|F(z)| \leq 1$ for all $z \in \mathbb{D}$ and $F(0) = |J_F(0)| - 1 = 0$, then $L(F)$ is univalent and maps $\mathbb{D}$ onto $\mathbb{D}$.

Proof. Let

$$H(z) := L(F) = zF_z(z) - \overline{z}F_{\overline{z}}(z) = \sum_{k=1}^{p}|z|^{2(k-1)}\Big(zh_k'(z) - \overline{z}\overline{g_k'(z)}\Big).$$

Then $H_z(z)$ and $H_{\overline{z}}(z)$ can be written as

$$\begin{aligned} H_z(z) &= h_1'(z) + zh_1''(z) + \sum_{k=2}^{p}\Big(k|z|^{2(k-1)}h_k'(z) + z|z|^{2(k-1)}h_k''(z) \\ &\quad -(k-1)\overline{z^2}|z|^{2(k-2)}\overline{g_k'(z)}\Big), \end{aligned}$$

and

$$\begin{aligned} -H_{\overline{z}}(z) &= \overline{g_1'(z)} + \overline{z}\overline{g_1''(z)} + \sum_{k=2}^{p}\Big(k|z|^{2(k-1)}\overline{g_k'(z)} + \overline{z}|z|^{2(k-1)}\overline{g_k''(z)} \\ &\quad -(k-1)z^2|z|^{2(k-2)}h_k'(z)\Big). \end{aligned}$$

Let r be a constant in $(0,1)$. For any distinct z_1, $z_2 \in \mathbb{D}_r$, we have

$$\begin{aligned} \big|H(z_1) - H(z_2)\big| &= \left|\int_{[z_1,z_2]} H_z(z)dz + H_{\overline{z}}(z)d\overline{z}\right| \\ &\geq \left|\int_{[z_1,z_2]} H_z(0)dz + H_{\overline{z}}(0)d\overline{z}\right| \\ &\quad - \left|\int_{[z_1,z_2]} \big(H_z(z) - H_z(0)\big)dz + \big(H_{\overline{z}}(z) - H_{\overline{z}}(0)\big)d\overline{z}\right| \\ &\geq K_1 - K_2 - K_3 - K_4, \end{aligned}$$

where

$$\begin{aligned}
K_1 &:= \left|\int_{[z_1,z_2]} h_1'(0)\,dz - \overline{g_1'(0)}d\overline{z}\right|,\\
K_2 &:= \left|\int_{[z_1,z_2]} \big(h_1'(z)-h_1'(0)\big)dz - \big(\overline{g_1'(z)}-\overline{g_1'(0)}\big)d\overline{z}\right|,\\
K_3 &:= \left|\int_{[z_1,z_2]} \sum_{k=1}^{p}|z|^{2(k-1)}\Big(zh_k''(z)dz - \overline{z}\overline{g_k''(z)}d\overline{z}\Big)\right|,\\
K_4 &:= \left|\int_{[z_1,z_2]} \sum_{k=2}^{p}(k-1)|z|^{2(k-2)}\Big(z^2h_k'(z)d\overline{z} - \overline{z^2}\overline{g_k'(z)}dz\Big)\right|\\
&\quad+\left|\int_{[z_1,z_2]} \sum_{k=2}^{p}k|z|^{2(k-1)}\Big(h_k'(z)dz - \overline{g_k'(z)}d\overline{z}\Big)\right|.
\end{aligned}$$

First, observe the following estimate for K_1:

$$K_1 \geq \int_{[z_1,z_2]} \lambda_F(0)|dz| = \lambda_F(0)|z_1-z_2|.$$

Next, by Theorem 1 (II), we have $|a_{n,k}|+|b_{n,k}| \leq \sqrt{M^4-1}\cdot\lambda_F(0)$ for $(n,k)\neq(1,1)$. Hence

$$\begin{aligned}
K_2 &= \left|\int_{[z_1,z_2]} \sum_{n=2}^{\infty} na_{n,1}z^{n-1}dz - \sum_{n=2}^{\infty} n\overline{b_{n,1}z^{n-1}}d\overline{z}\right|\\
&\leq |z_1-z_2|\sum_{n=2}^{\infty} n(|a_{n,1}|+|b_{n,1}|)r^{n-1}\\
&\leq |z_1-z_2|\sqrt{M^4-1}\cdot\lambda_F(0)\frac{2r-r^2}{(1-r)^2},
\end{aligned}$$

and

$$\begin{aligned}
K_3 &\leq |z_1-z_2|\sum_{k=1}^{p} r^{2(k-1)}\sum_{n=2}^{\infty} n(n-1)(|a_{n,k}|+|b_{n,k}|)r^{n-1}\\
&\leq |z_1-z_2|\sqrt{M^4-1}\cdot\lambda_F(0)\sum_{k=1}^{p}\frac{2r^{2k-1}}{(1-r)^3}.
\end{aligned}$$

Finally,

$$\begin{aligned} K_4 &\leq \int_{[z_1,z_2]} \sum_{k=2}^{p}(k-1)r^{2(k-2)}\sum_{n=1}^{\infty} n(|a_{n,k}|+|b_{n,k}|)r^{n+1}|dz| \\ &\quad + \int_{[z_1,z_2]} \sum_{k=2}^{p} kr^{2(k-1)}\sum_{n=1}^{\infty} n(|a_{n,k}|+|b_{n,k}|)r^{n-1}|dz| \\ &\leq |z_1-z_2|\sum_{k=2}^{p}(2k-1)r^{2(k-1)}\sum_{n=1}^{\infty} n(|a_{n,k}|+|b_{n,k}|)r^{n-1} \\ &\leq |z_1-z_2|\sqrt{M^4-1}\cdot\lambda_F(0)\sum_{k=2}^{p}(2k-1)\frac{r^{2(k-1)}}{(1-r)^2}. \end{aligned}$$

Using these estimates, we obtain

$$\big|H(z_1)-H(z_2)\big| \geq K_1-K_2-K_3-K_4 \geq \lambda_F(0)|z_1-z_2|\varphi(r),$$

where

$$\varphi(r) = 1-\sqrt{M^4-1}\left(\frac{2r-r^2}{(1-r)^2}+\sum_{k=1}^{p}\frac{2r^{2k-1}}{(1-r)^3}+\sum_{k=2}^{p}\frac{(2k-1)r^{2(k-1)}}{(1-r)^2}\right).$$

It is easy to verify that $\varphi(r)$ is a strictly decreasing function for $r\in(0,1)$ and $M>1$,

$$\lim_{r\to 0+}\varphi(r)=1 \text{ and } \lim_{r\to 1^-}\varphi(r)=-\infty.$$

Hence there exists a unique $r_6\in(0,1)$ satisfying $\varphi(r_6)=0$. This shows that $H(z)$ is univalent in $\mathbb{D}_{r_6}$.

Furthermore, for any w in $\{w:\ |w|=r_6\}$,

$$\begin{aligned} \big|H(w)\big| &= \left|\sum_{k=1}^{p}|w|^{2(k-1)}\big(wh_k'(w)-\overline{w}\overline{g_k'(z)}\big)\right| \\ &\geq \big|wh_1'(0)-\overline{w}\overline{g_1'(0)}\big| - \Big|w\big(h_1'(w)-h_1'(0)\big)-\overline{w}\big(\overline{g_1'(w)}-\overline{g_1'(0)}\big)\Big| \\ &\quad -\left|\sum_{k=2}^{p}|w|^{2(k-1)}\big(wh_k'(w)-\overline{w}\overline{g_k'(w)}\big)\right| \\ &\geq \lambda_F(0)r_6-\sum_{n=2}^{\infty}n(|a_{n,1}|+|b_{n,1}|)r_6^n-\sum_{k=2}^{p}r_6^{2(k-1)}\sum_{n=1}^{\infty}n(|a_{n,k}|+|b_{n,k}|)r_6^n. \end{aligned}$$

By Theorem 1 (II), we see $|a_{n,k}|+|b_{n,k}|\leq\sqrt{M^4-1}\cdot\lambda_F(0)$ for all $(n,k)\neq(1,1)$. Therefore, we obtain

$$\begin{aligned}\big|H(w)\big| &\geq \lambda_F(0)r_6\left(1-\sqrt{M^4-1}\frac{2r_6-r_6^2}{(1-r_6)^2}-\sqrt{M^4-1}\sum_{k=2}^{p}\frac{r_6^{2(k-1)}}{(1-r_6)^2}\right)\\ &\geq \lambda_0(M)r_6\left(1-\sqrt{M^4-1}\frac{2r_6-r_6^2}{(1-r_6)^2}-\sqrt{M^4-1}\sum_{k=2}^{p}\frac{r_6^{2(k-1)}}{(1-r_6)^2}\right):=\rho_6,\end{aligned}$$

where $\lambda_0(M)$ is the same as in (6), and

$$\begin{aligned}\rho_6 &> \lambda_0(M)r_6\left[1-\sqrt{M^4-1}\left(\frac{2r_6-r_6^2}{(1-r_6)^2}+\sum_{k=1}^{p}\frac{2r_6^{2k-1}}{(1-r_6)^3}\right.\right.\\ &\qquad\left.\left.+\sum_{k=2}^{p}\frac{(2k-1)r_6^{2(k-1)}}{(1-r_6)^2}\right)\right]=0.\end{aligned}$$

If $|F(z)|\leq 1$ for all $z\in\mathbb{D}$ and $F(0)=|J_F(0)|-1=0$, then by Corollary 2 (I), we have $L(F)$ is univalent and maps $\mathbb{D}$ onto $\mathbb{D}$. The proof of the theorem is complete. □

Corollary 5. *Suppose that F is a polyharmonic mapping of the form* (2) *and all its non-zero coefficients a_{n,k_1}, a_{n,k_2} and b_{n,k_3}, b_{n,k_4} satisfy* (3).

If $|F(z)|\leq M$ in $\mathbb{D}$ for some $M>1$ and $F(0)=\lambda_F(0)-1=0$, then $L(F)$ is univalent in the disk $\mathbb{D}_{r_7}$ and $F(\mathbb{D}_{r_7})$ contains a univalent disk $\mathbb{D}_{\rho_7}$, where $r_7(M,p)$ is the least positive root of the following equation:

$$1-\sqrt{2M^2-2}\left(\frac{2r-r^2}{(1-r)^2}+\sum_{k=1}^{p}\frac{2r^{2k-1}}{(1-r)^3}+\sum_{k=2}^{p}\frac{(2k-1)r^{2(k-1)}}{(1-r)^2}\right)=0,$$

and

$$\rho_7=r_7\left(1-\sqrt{2M^2-2}\frac{2r_7-r_7^2}{(1-r_7)^2}-\sqrt{2M^2-2}\sum_{k=2}^{p}\frac{r_7^{2(k-1)}}{(1-r_7)^2}\right).$$

In particular, if $|F(z)|\leq 1$ for all $z\in\mathbb{D}$ and $F(0)=\lambda_F(0)-1=0$, then $L(F)$ is univalent and maps $\mathbb{D}$ onto $\mathbb{D}$.

Example 3. *Under the hypothesis of Example 2, and by Corollary 5, $L(F_1)$ is univalent in the disk $\mathbb{D}_{r_8}$ and $F_1(\mathbb{D}_{r_8})$ contains a univalent disk $\mathbb{D}_{\rho_8}$, where $r_8\approx$ 0.00798 is the least positive root of the following equation:*

$$1-\sqrt{2M_1^2-2}\cdot\frac{4r-3r^2+3r^3+3r^4-3r^5}{(1-r)^3}=0,$$

and

$$\rho_8=r_8\left(1-\sqrt{2M_1^2-2}\cdot\frac{2r_8}{(1-r_8)^2}\right)\approx 0.00400,$$

while by [5, Theorem 1.1]*, we see that* $L(F_1)$ *is univalent in the disk* $\mathbb{D}_{r_9}$ *and* $F(\mathbb{D}_{r_9})$ *contains a univalent disk* $\mathbb{D}_{\rho_9}$*, where* $r_9 \approx 0.00013$ *is the least positive root of the following equation:*

$$\frac{\pi}{4M_2} - \frac{6M_2 r^2}{(1-r)^2} - \frac{4Mr^3}{(1-r)^3} - \frac{16M_2}{\pi^2} m_1 \arctan r - \frac{4M_2 r}{(1-r)^3} = 0,$$

where $m_1 \approx 6.05934$ *is the minimum value of the function*

$$\frac{2 - x^2 + \frac{4}{\pi}\arctan x}{x(1-x^2)}$$

for $0 < x < 1$*, and*

$$\rho_9 = r_9 \left(\frac{\pi}{4M_2} - \frac{2M_2 r_9^2}{(1-r_9)^2} - \frac{16M_2}{\pi^2} m_1 \arctan r_9 \right) \approx 1.48687 \times 10^{-6}.$$

References

1. Z. Abdulhadi and Y. Abu Muhanna, Landau's theorem for biharmonic mappings. *J. Math. Anal. Appl.* **338** (2008), 705–709.
2. Z. Abdulhadi, Y. Abu Muhanna and S. Khuri, On univalent solutions of the biharmonic equation. *J. Inequal. Appl.* **5** (2005), 469–478.
3. Z. Abdulhadi, Y. Abu Muhanna and S. Khuri, On some properties of solutions of the biharmonic equation. *Appl. Math. Comput.* **117** (2006), 346–351.
4. H. Chen, P. M. Gauthier and W. Hengartner, Bloch constants for planar harmonic mappings. *Proc. Amer. Math. Soc.* **128** (2000), 3231–3240.
5. Sh. Chen, S. Ponnusamy and X. Wang, Landau's theorem for certain biharmonic mappings. *Appl. Math. Comput.* **208** (2009), 427–433.
6. Sh. Chen, S. Ponnusamy and X. Wang, Properties of some classes of planar harmonic and planar biharmonic mappings. *Complex Anal. Oper. Theory* **5** (2011), 901–916.
7. Sh. Chen, S. Ponnusamy and X. Wang, Coefficient estimates, Landau's theorem and Bohr's inequality for harmonic mappings. Submitted.
8. Sh. Chen, S. Ponnusamy and X. Wang, Bloch constant and Landau's theorem for planar p-harmonic mappings. *J. Math. Anal. Appl.* **373** (2011), 102–110.
9. J. Chen, A. Rasila and X. Wang, On polyharmonic univalent mappings. *Preprint. arXiv:1302.2018.*
10. J. Chen, A. Rasila and X. Wang, Starlikeness and convexity of polyharmonic mappings. *Preprint. arXiv:1302.2398.*
11. J. Chen and X. Wang, On certain classes of biharmonic mappings defined by convolution. *Abstr. Appl. Anal.* **2012**, Article ID 379130, 10 pages. doi:10.1155/2012/379130
12. J. G. Clunie and T. Sheil-Small, Harmonic univalent functions. *Ann. Acad. Sci. Fenn. Ser. A. I.* **9** (1984), 3–25.
13. M. Dorff and M. Nowak, Landau's theorem for planar harmonic mappings. *Comput. Methods Funct. Theory* **4** (2004), 151–158.
14. P. Duren, *Harmonic mappings in the plane.* Cambridge University Press, Cambridge, 2004.
15. A. Grigoryan, Landau and Bloch theorems for planar harmonic mappings. *Complex Var. Elliptic Equ.* **51** (2006), 81–87.
16. J. Happel and H. Brenner, *Low Reynolds number hydrodynamics with special applications to particulate media.* Prentice-Hall, Englewood Cliffs, NJ, USA, 1965.
17. S. A. Khuri, Biorthogonal series solution of Stokes flow problems in sectorial regions. *SIAM J. Appl. Math.* **56** (1996), 19–39.
18. W. E. Langlois, *Slow viscous flow.* Macmillan, New York, NY, USA, 1964.

19. M. Liu, Landau theorems for biharmonic mappings. *Complex Var. Elliptic Equ.* **53** (2008), 843–855.
20. Y. Zhu and M. Liu, Landau-type theorems for certain planar harmonic mappings or biharmonic mappings. *Complex Var. Elliptic Equ.* (2012), 1–10 iFirst.
21. M. Liu, Estimates on Bloch constants for planar harmonic mappings. *Sci. China Ser. A* **52** (2009), 87–93.
22. J. Qiao and X. Wang, On p-harmonic univalent mappings (in Chinese). *Acta Math. Sci.* **32A** (2012), 588–600.

J. Chen, Department of Mathematics, Hunan Normal University, Changsha, Hunan 410081, People's Republic of China.
E-mail address: `jiaolongchen@sina.com.cn`

A. Rasila, Department of Mathematics, Hunan Normal University, Changsha, Hunan 410081, People's Republic of China, and Department of Mathematics and Systems Analysis, Aalto University, P. O. Box 11100, FI-00076 Aalto, Finland.
E-mail address: `antti.rasila@iki.fi`

X. Wang, Department of Mathematics, Hunan Normal University, Changsha, Hunan 410081, People's Republic of China.
E-mail address: `xtwang@hunnu.edu.cn`